

# Nonlinear modes for the Gross-Pitaevskii equation – demonstrative computation approach

G L Alfimov and D A Zezyulin

Moscow Institute of Electronic Engineering, Zelenograd, Moscow, 124498, Russia

E-mail: galfimov@yahoo.com, d.zezyulin@gmail.com

**Abstract.** A method for the study of steady-state nonlinear modes for Gross-Pitaevskii equation (GPE) is described. It is based on exact statement about coding of the steady-state solutions of GPE which vanish as  $x \rightarrow +\infty$  by reals. This allows to fulfill *demonstrative computation* of nonlinear modes of GPE i.e. the computation which allows to guarantee that *all* nonlinear modes within a given range of parameters have been found. The method has been applied to GPE with quadratic and double-well potential, for both, repulsive and attractive nonlinearities. The bifurcation diagrams of nonlinear modes in these cases are represented. The stability of these modes has been discussed.

PACS numbers: 03.75.Hh, 02.70.-c, 02.30.Oz

Submitted to: *Nonlinearity*

## 1. Introduction

The Gross-Pitaevskii equation (GPE) is one of the basic models in the theory of Bose-Einstein condensate (BEC)[1]. In dimensionless form this equation reads

$$i\Psi_t = \Delta\Psi - V(\mathbf{x})\Psi + \sigma\Psi|\Psi|^2, \quad (1)$$

In BEC context GPE describes an atomic cloud in the mean-field approximation. The function  $\Psi(t, \mathbf{x})$  corresponds to macroscopic wave function,  $\Delta$  is the Laplace operator and  $V(\mathbf{x})$  is a trap potential for the condensate to be confined. The value of  $\sigma$  governs the sign of nonlinearity: the case  $\sigma = 1$  corresponds to attractive interatomic interactions and  $\sigma = -1$  corresponds to repulsive interatomic interactions. Both the cases  $\sigma = \pm 1$  are of interest for physical applications [2, 3]. Eq.(1) conserves the quantities

$$E = \frac{1}{2} \int \left( |\nabla\Psi|^2 - V(\mathbf{x})|\Psi|^2 - \frac{\sigma}{2}|\Psi|^4 \right) d\mathbf{x} \quad (2)$$

$$N = \int |\Psi|^2 d\mathbf{x} \quad (3)$$

which have the sense of Hamiltonian and number of particles correspondingly. An important class of solutions of GPE are stationary nonlinear modes defined as

$$\Psi(t, \mathbf{x}) = e^{i\omega t}\psi(\mathbf{x}) \quad (4)$$

$$\psi(\mathbf{x}) \rightarrow 0 \quad \text{as} \quad |\mathbf{x}| \rightarrow \infty \quad (5)$$

In this case  $\psi(\mathbf{x})$  satisfies the equation

$$\Delta\psi + (\omega - V(\mathbf{x}))\psi + \sigma\psi|\psi|^2 = 0, \quad (6)$$

The parameter  $\omega$  in terms of BEC corresponds to the chemical potential. The *ground state* solutions of (5)-(6) (i.e. node-free solutions which minimize energy functional for Eq.(1)) are of primary importance for BEC applications [4, 6, 5]. Apart from them some *non-ground* (i.e. excited) nonlinear modes also have been reported (see e.g. [7, 9, 11, 12, 13]) as well as physical mechanisms for their generation [8]. Except some specific examples of the potential  $V(\mathbf{x})$  when exact solutions of (5)-(6) are available (see e.g. [14]) in all the cases the nonlinear states have been found numerically.

Typically, numerical calculation of nonlinear modes needs some initial guess for subsequent iterative procedure. In order to get *a priori* information about the shape of nonlinear states various approaches have been used. In [9] the branches of nonlinear modes for the case  $V(\mathbf{x}) = |\mathbf{x}|^2$  have been found by continuation from the limit  $\psi \rightarrow 0$  where the nonlinear term can be neglected

$$\Delta\psi + (\omega - |\mathbf{x}|^2)\psi = 0. \quad (7)$$

Therefore, the shapes of these states can be described by the set of Gauss-Hermite functions. In 1D case and for  $\sigma = 1$  it has been proved rigorously [10] that there are infinitely many branches of nonlinear modes which bifurcate from zero solution of (6) at discrete infinite set of values of  $\omega$  corresponding to eigenvalues of linear problem (7). Following [11, 12], it is quite natural to call these modes *the modes with linear*

*counterpart*. It has been shown in [11, 12] that for some potentials  $V(\mathbf{x})$  nonlinear modes without linear counterpart also can exist. In the case of multi-well potentials these states have been found by continuation from asymptotical limit of large nonlinearity [11, 12].

However, the following question remains: can one list *all* nonlinear states corresponding to given potential  $V(\mathbf{x})$  with  $\omega$  lying within some range? To best of our knowledge, up to the moment there is no answer to this question even in the most explored 1D-case of quadratic potential  $V(x) = x^2$ . The aim of this paper is to make a step toward answering this question. Specifically, for  $\sigma = -1$  we explain how to arrange numerical calculation in such a way that *all* nonlinear modes described by Eq.(6) for  $\omega$  lying within a given range would be found. In the case  $\sigma = 1$  the method gives a picture which is complete, as we believe, but this fact has not been proved. We illustrate our approach on 1D case of Eq.(6) with quadratic and double-well potentials (the applications to radial 2D and 3D cases are straightforward). The base for this approach is given by *exact* statement on one-to-one correspondence between the solutions of Eq.(6) which tends to zero as  $x \rightarrow +\infty$  and linear coefficient  $C_+$  which determines their asymptotics in this limit.

So, in this paper we consider 1D version of Eq.(6), specifically

$$\psi'' + (\omega - V(x))\psi + \sigma\psi^3 = 0 \quad (8)$$

where prime denotes differentiation with respect to  $x$ . We assume that the potential  $V(x)$  is *even*,  $V(x) = V(-x)$  and grows when  $|x| \rightarrow \infty$ . The attention is focused on two particular examples of the potential,

$$V(x) = x^2, \quad (\text{quadratic potential}) \quad (9)$$

$$V(x) = \alpha x^4 + \beta x^2, \quad \alpha > 0, \quad \beta < 0 \quad (\text{double-well potential}) \quad (10)$$

The solutions of Eq.(8) are supposed to be real and satisfy the localization condition

$$\lim_{x \rightarrow \pm\infty} \psi(x) = 0 \quad (11)$$

The paper is organized as follows. In Section 2 we describe the method which we use. In Section 3 we apply this method to Eq.(8) with quadratic and double-well potentials. Section 4 contains summary and discussion. Some auxiliary statements and technical details can be found in Appendices A-C.

## 2. The description of the approach

### 2.1. The idea of the approach

The approach which we propose is an extension of one used in [6, 20] for numerical construction of nonlinear modes for Eq.(6). Let us denote by  $S_+$  the set of all solutions of Eq.(8) which satisfy the condition

$$\lim_{x \rightarrow +\infty} \psi(x) = 0 \quad (12)$$

When  $x \rightarrow +\infty$  the nonlinear term in Eq.(8) can be neglected and the asymptotics of solution from  $S_+$  can be found from linearized equation

$$\psi'' + (\omega - V(x))\psi = 0 \quad (13)$$

The theory of Eq.(13) (see e.g. [15, 16]) says that for a wide class of potentials  $V(x)$  it has *unique* (up to constant factor) solution  $\tilde{\psi}(x)$  such that  $\tilde{\psi}(x) \rightarrow 0$  when  $x \rightarrow +\infty$ . If  $V(x) > \omega$  for  $x > x_0$  for some  $x_0$  large enough (and fixed) then the asymptotics of this specific solution is described by the formula [16]

$$\tilde{\psi}(x) = \frac{e^{-\int_{x_0}^x \sqrt{V(t)-\omega} dt}}{\sqrt[4]{V(x)-\omega}} (1 + o(1)), \quad x \rightarrow +\infty \quad (14)$$

Then the solution of Eq.(8),  $\psi(x)$ , which satisfies (12) has the asymptotics

$$\psi(x) = \frac{e^{-\int_{x_0}^x \sqrt{V(t)-\omega} dt}}{\sqrt[4]{V(x)-\omega}} (C_+ + o(1)), \quad x \rightarrow +\infty \quad (15)$$

Here  $C_+$  is some constant. The next step is to assume that there is one-to-one correspondence between the solutions from  $S_+$  and the constants  $C_+ \in \mathbf{R}$  which comes from formula (15) (this fact will be rigorously justified below). This implies that the solutions  $S_+$  constitute one parameter set with free parameter  $C_+ \in \mathbf{R}$ . Let us denote a solution of Eq.(8) which satisfies the asymptotical relation (15) by  $\psi_+(x, C_+)$ . By construction, the solutions  $\psi_+(x; C_+)$  tend to zero as  $x \rightarrow +\infty$ ; however, generically, they do not tend to zero as  $x \rightarrow -\infty$ .

The same reasoning holds for the solutions of Eq.(8) which tend to zero as  $x \rightarrow -\infty$  (the set of these solutions will be denoted by  $S_-$ ). The asymptotics of these solutions is of the form

$$\psi(x) = \frac{e^{\int_x^{-x_0} \sqrt{V(t)-\omega} dt}}{\sqrt[4]{V(x)-\omega}} (C_- + o(1)), \quad x \rightarrow -\infty \quad (16)$$

where  $C_-$  is some constant. Assume that  $S_-$  is one-parameter set with free parameter  $C_-$ . Let us denote a solution of Eq.(8) which satisfies the asymptotical relation (16) by  $\psi_-(x, C_-)$ . Evidently, a localized solution  $\psi(x)$  which satisfies (11) must belong to both sets,  $S_+$  and  $S_-$ , and in the point  $x = 0$  the relations must be valid

$$\psi(0) = \psi_-(0, C_-) = \psi_+(0, C_+); \quad \psi'(0) = \psi'_-(0, C_-) = \psi'_+(0, C_+); \quad (17)$$

In fact, (17) is a system of equations to determine unknown constants  $C_-$  and  $C_+$  and, consequently, a solution  $\psi(x)$ . In practice, the values  $\psi_+(0; C_+)$  and  $\psi'_+(0, C_+)$  can be calculated numerically (using, for example, Runge-Kutta method) by solving the initial value problem for Eq.(8) starting from some large enough value  $x = x_\infty$  where the values  $\psi_+(x_\infty; C_+)$ ,  $\psi'_+(x_\infty; C_+)$  can be approximated by means of asymptotical formula (15). In the same way the values  $\psi_-(0; C_-)$ ,  $\psi'_-(0; C_-)$  can be calculated.

## 2.2. Mathematical basis

The heuristical arguments given in paragraph 2.1 can be justified by exact statement which describes the set of solutions of Eq.(8) which tend to zero as  $x \rightarrow +\infty$ . This can be done using “asymptotic integration” of Eq.(8) (see e.g. [19]). The following statement is valid:

Theorem 1. *Let there exists some  $x_0$  such that*

$$(i) \ V(x) - \omega \geq \varepsilon > 0 \text{ for } x > x_0;$$

$$(ii) \ V(x) \in C^2(x_0, +\infty);$$

$$(iii) \ \int_{x_0}^{\infty} \frac{|V''(x)|}{(V(x)-\omega)^{3/2}} dx < \infty$$

*Let  $\psi_1(x)$  be a fixed solution of linear equation (13) such that  $\psi_1(x) \rightarrow 0$  as  $x \rightarrow +\infty$  which has asymptotics (14). Then*

*(a) for any solution  $\psi(x) \in S_+$  there exists a limit*

$$\lim_{x \rightarrow +\infty} \frac{\psi(x)}{\psi_1(x)} = C_+ < \infty \quad (18)$$

*(b) for any  $C_+ \in \mathbf{R}$  there exists an unique solution  $\psi_+(x; C_+) \in S_+$  of Eq.(8) such that*

$$\lim_{x \rightarrow +\infty} \frac{\psi_+(x; C_+)}{\psi_1(x)} = C_+ \quad (19)$$

*(c) if  $\psi_+(x; C_+)$  exists in some neighbourhood  $U$  of a point  $(\tilde{x}; \tilde{C}_+)$  then in  $U$  there also exists partial derivative  $\frac{\partial \psi_+(x; C_+)}{\partial C_+}$  which is continuous with respect to  $C_+$ .*

Theorem 1 establishes one-to-one correspondence between  $S_+$  and  $\mathbf{R}$ ; specifically, it says that the set  $S_+$  can be parametrized by real parameter  $C_+ \in \mathbf{R}$ . The point (a) of Theorem 1 has been used for the search of localized solutions by means of shooting method (see e.g. [20]). We make use of the point (b) to construct *general picture* of branches of solutions of Eq.(8). The point (c) will be used for proving Lemma 1 in Appendix C.

Theorem 1 reflects the well-known relation between the behavior of linear dynamical system and its nonlinear perturbation in a vicinity of its zero solution. In spite of bulk of literature devoted to this subject (see, for instance, [17] and references therein) we failed to find this statement *exactly* in the form we need. Therefore we adduce a sketch of proof of Theorem 1 in Appendix B in a little more general form. Evidently, analogous statement can be made for the solutions of Eq.(8) which tend to zero as  $x \rightarrow -\infty$ .

## 2.3. Practical implementation of the method and visualization of the results

**2.3.1.  $(\psi(0), \psi'(0))$ -pictures.** Let the potential  $V(x)$  be *even*,  $V(x) = V(-x)$ . Let  $\omega$  be fixed and  $x = x_\infty$  is some value large enough where the solution of (8) can be approximated well by the asymptotics (15). Assume that the numerical continuation of the solutions  $\psi_+(x; C_+)$  from  $x = x_\infty$  to  $x = 0$  can be fulfilled for each  $C_+$  in some

interval  $C_+ \in [-\tilde{C}_+; \tilde{C}_+]$  (using, for instance, Runge-Kutta method). The result can be represented as a segment of curve

$$\gamma_+ : \{(\psi_+(0; C_+); \psi'_+(0; C_+)), C_+ \in [-\tilde{C}_+; \tilde{C}_+]\}$$

on the plane  $(\psi; \psi')$ . Since  $C_+ = 0$  corresponds to zero solution, this curve passes through the point  $(0; 0)$  of this plane. Moreover, due to the fact that the equation (8) admits the symmetry  $\psi \rightarrow -\psi$ , this segment of the curve is centrosymmetric with respect to the origin. In practice this means that the calculation should be fulfilled for  $[0; \tilde{C}_+]$  only. In analogous way, let us consider the segment of curve

$$\gamma_- : \{(\psi_-(0; C_-); \psi'_-(0; C_-)), C_- \in [-\tilde{C}_-; \tilde{C}_-]\}$$

constructed by means of numerical continuation of the solution  $\psi_-(x; C_-)$  from the value  $x = -x_\infty$  until the value  $x = 0$ . Since equation (8) is invariant with respect to the change  $x \rightarrow -x$ , if  $\tilde{C}_- = \tilde{C}_+$  the curve  $\gamma_-$  is symmetric about the  $\psi'$  axis to the curve  $\gamma_+$ . Then, each point of intersection of  $\gamma_-$  and  $\gamma_+$  corresponds to a solution of system (17) and, consequently, to a solution of Eq.(8) with the condition (11) (see Fig.1).

Due to the symmetries of the curves  $\gamma_-$  and  $\gamma_+$ , the points of intersection of  $\gamma_+$  with the axes  $\psi$  and  $\psi'$  are also points of intersection of  $\gamma_+$  with the curve  $\gamma_-$ . These points correspond to specific localized solutions of Eq.(8); specifically, the points of intersection of  $\gamma_+$  with the axis  $\psi$  correspond to *even* localized solutions of Eq.(8) whereas the points of intersection of  $\gamma_+$  with the axis  $\psi'$  correspond to *odd* localized solutions of this equation. The points of intersection of  $\gamma_-$  and  $\gamma_+$  which do not belong to the axes correspond to non-symmetric localized solutions of Eq.(8).

**2.3.2.  $(C_+, \omega)$ -diagrams.** Let us consider now branches of localized solutions of Eq.(8) which appear when the parameter  $\omega$  varies. Due to Theorem 1, if  $\omega$  is fixed, each of the localized solutions of Eq.(8) can be unambiguously defined by the value  $C_+$  determined by its asymptotics. Therefore, it is convenient to represent the branches of localized solutions on the plane  $(C_+, \omega)$  marking for each  $\omega$  those values  $C_+$  for which localized solutions exist. In such a way any branch of solutions corresponds to a curve  $\Gamma$  on the plane  $(C_+, \omega)$ . An advantage of this graphical representation (in comparison, for instance, with more traditional  $(N, \omega)$  representation where  $N$  is given by (3)) is that *each point of the plane  $(C_+, \omega)$  corresponds to at most one localized solution* of Eq.(8). This means, for instance, that if two curves, corresponding to two branches have a common point, then there exists a localized solution which belongs to both branches. This is not the case for  $(N, \omega)$ -diagrams.

Consider now any segment of curve  $\Gamma$  which has no intersection with the axis  $C_+ = 0$ . Simple reasoning shows that *the number of zeros of solutions  $\psi(x)$  is constant along this segment of  $\Gamma$* . Indeed, consider a continuous deformation of solution  $\psi(x)$  which occurs when passing along this segment. A zero  $x = x_0$  of  $\psi(x)$  can disappear by (i) merging with one or several neighboring zeros; (ii) by merging with  $x = \pm\infty$  ("goes to infinity"). The case (i) implies that the branch includes a solution with  $\psi(x_0) = \psi'(x_0) = 0$  which is zero solution. Consider the case (ii) i.e. assume that zero

$x = x_0$  merges with  $x = +\infty$ . Then for some solution of the branch the parameter  $C_+$  is zero which implies that the solution is identically zero. Similar arguments allow to discard also the case when (i) and (ii) hold simultaneously.

#### 2.4. Some additional statements for the case of repulsive nonlinearity

In the case of repulsive nonlinearity Eq.(8) reads

$$\psi'' + (\omega - V(x))\psi - \psi^3 = 0 \quad (20)$$

It turns out that “most part” of solutions of this equation *are singular* i.e. they tend to  $\pm\infty$  as  $x$  approaches some finite value. This fact allows to fulfill more complete analysis of localized solutions for Eq.(20). It can be based on the following statement:

**Lemma 1** *Let there exists a solution  $\psi_0(x) = \psi_+(x, C_+^0) \in S_+$ ,  $C_+^0 > 0$  of equation (20) such that:*

(i)  $\psi_0(x)$  is defined on  $(x_0; +\infty)$  where  $x_0 > 0$  and

$$\lim_{x \rightarrow x_0+0} \psi_0(x) = +\infty$$

(ii)  $\psi_0(x) > 0$  for  $x \in (x_0; +\infty)$ ;

(iii) for  $x \in (x_0; +\infty)$  the following inequality holds:

$$\omega - V(x) - 3\psi_0^2(x) < 0$$

*Then any solution  $\psi_+(x, C_+) \in S_+$  for  $C_+ > C_+^0$  has a singularity at some point  $x = x(C_+) > x_0$ ,  $\lim_{x \rightarrow x(C_+)+0} \psi_+(x, C_+) = +\infty$ .*

The proof of Lemma 1 is postponed in Appendix C. Lemma 1 allows to stop the scanning  $C_+$  when  $C_+^0$  satisfying the conditions of Lemma 1 is achieved. The window of scanning in this case is *finite*, so one can conclude that *all* the solutions which exist for given  $\omega$  have been found. However, for practical implementation of Lemma 1 in numerics one needs a condition to assure that the solution which has been found is singular.

Let us introduce the value

$$\Omega \equiv \sup_{x \in \mathbf{R}} (\omega - V(x))$$

and assume that  $|\Omega| < \infty$ . If  $\Omega \leq 0$  then one can prove that Eq.(20) has no localized solutions. If  $\Omega > 0$  the following statement holds

**Lemma 2** *Let  $\Omega > 0$ . If a solution  $\psi(x)$  of Eq.(20) satisfies at some point  $x = x_0$  the condition  $|\psi(x_0)| > \sqrt{\Omega}$  then this solution is singular.*

The proof of Lemma 2 is also postponed in Appendix C. Lemma 2 allows to stop the shooting procedure once the value  $\psi = \sqrt{\Omega}$  is achieved.




**Figure 1.**  $(\psi(0), \psi'(0))$ -pictures for Eq.(8) with  $V(x) = x^2$ . Left panel: repulsive nonlinearity,  $\omega = 4$ ; right panel: attractive nonlinearity,  $\omega = 0$ .

### 3. Numerical results.

#### 3.1. The quadratic potential.

**3.1.1. Repulsive nonlinearity** First, we applied the approach described above for the equation

$$\psi'' + (\omega - x^2)\psi - \psi^3 = 0 \quad (21)$$

taking  $0 \leq \omega \leq 12$ . According to the formula (14) the asymptotics of the solution  $\psi(x)$  as  $x \rightarrow +\infty$  is

$$\psi(x) \sim C_+ x^{\frac{1}{2}(\omega-1)} e^{-\frac{x^2}{2}} \quad (22)$$

The starting point for the shooting,  $x_\infty$ , was taken equal to 10. The results were recalculated for  $x_\infty = 12$  and no significant change was observed. The zoom of scanning with respect to  $C_\pm$  was from  $C_\pm = 0$  until the value  $C_\pm^{(0)}$  where the conditions of Lemma 1 held (the value  $C_\pm^{(0)}$  has been found for all considered  $\omega$ ). This guarantees that *all* localized solutions within this range of  $\omega$  were found.

The Fig.1, left panel, shows the  $(\psi(0), \psi'(0))$ - picture for  $\omega = 4$ . In Fig.2(a)-(b) six branches of nonlinear modes for Eq.(8) with repulsive nonlinearity are depicted. All these branches,  $\Gamma_r^n$ , are originated by localized modes of linear harmonic oscillator, corresponding to eigenvalues  $\omega = 2n - 1$ ,  $n = 1 \div 6$ . The linear limit can be achieved by passing  $C_+ \rightarrow 0$  (or, correspondingly,  $N \rightarrow 0$ ). The branches of even and odd solutions alternate. It follows from these figures that there are *no other branches* of localized modes within this range of  $\omega$ . The analysis of linear stability for the branches  $\Gamma_r^n$  shows that the solutions corresponding to the branches  $\Gamma_r^1$  and  $\Gamma_r^2$  are stable whereas




**Figure 2.** The branches of nonlinear modes for Eq.(8) with  $V(x) = x^2$ . (a):  $(C_+, \omega)$ -diagrams, repulsive nonlinearity; (b):  $(N, \omega)$ -diagrams, repulsive nonlinearity; (c)  $(C_+, \omega)$ -diagrams, attractive nonlinearity; (d)  $(N, \omega)$ -diagrams, attractive nonlinearity. All the modes are originated by the modes of linear harmonic oscillator.

the solutions of the other two branches (and, probably, of all the higher branches) are unstable. This completely agrees with results of [21], (see Fig. 1 of [21]).

*3.1.2. Attractive nonlinearity* For the case of attractive nonlinearity,

$$\psi'' + (\omega - x^2)\psi + \psi^3 = 0 \quad (23)$$

it is proved [10] that each localized mode of harmonic oscillator, corresponding to  $\omega = 2n - 1$ ,  $n = 1, 2, \dots$  originates a branch of nonlinear modes,  $\Gamma_a^n$ . The asymptotics of the solution  $\psi(x)$  as  $x \rightarrow +\infty$  is described by the formula (22). We study Eq.(23) for  $-5 \leq \omega \leq 5$ . Since in this case Lemma 1 cannot be applied the scanning with respect to  $C_{\pm}$  was fulfilled within the fixed range  $0 \leq C_{\pm} \leq 10$ . Fig.1 (right panel) shows  $(\psi(0), \psi'(0))$ - picture for  $\omega = 0$ . Fig.2(c) represents  $(C_+, \omega)$ -diagram and Fig.2(d) shows the corresponding  $(N, \omega)$ -diagram. The four branches  $\Gamma_a^n$ ,  $n = 1, 2, 3, 4$  originated by the localized modes of harmonic oscillator have been found and no other modes have appeared within the considered range of  $C_{\pm}$  and  $\omega$ . We suppose that, in general, the branches of solutions generated by the modes of linear oscillator constitute *all* nonlinear

localized modes that can be described by Eq.(23).

Linear stability analysis shows that the solutions of the two lowest branches  $\Gamma_a^1$  and  $\Gamma_a^2$  are *stable* and the solutions of the branch  $\Gamma_a^4$  (and, probably, of all the higher branches  $\Gamma_a^n$ ,  $n \geq 4$ ) are unstable. This conforms with the results of [21], (see Fig.1 of [21]). At the same time we observed that the two-node solutions of the branch  $\Gamma_a^3$  are unstable in small-amplitude limit (for  $\omega^* < \omega < 5$ ,  $\omega^* \approx 3.83$ ) but they become stable for larger amplitudes ( $\omega < \omega^*$ ). Probably this is the reason of troubles with generation of this states by adiabatic variation of scattering length reported in [21].

### 3.2. The double-well potential

*3.2.1. Repulsive nonlinearity.* In this case the Eq.(8) reads

$$\psi'' + (\omega - \alpha x^4 - \beta x^2)\psi - \psi^3 = 0, \quad \alpha > 0, \quad \beta < 0 \quad (24)$$

According to the formula (14) the asymptotics of the solution  $\psi(x)$  as  $x \rightarrow +\infty$  is

$$\psi(x) \sim \frac{C_+}{x} \cdot \exp \left\{ -\frac{\sqrt{\alpha}}{3}x^3 - \frac{\beta}{2\sqrt{\alpha}}x - \frac{1}{2\sqrt{\alpha}} \left( \omega^2 + \frac{\beta^2}{4\alpha} \right) x^{-1} \right\} \quad (25)$$

All the calculations were done for the parameters of potential  $\alpha = 10$ ,  $\beta = -20$  for  $-5 \leq \omega \leq 10$ . The range of scanning for  $C_{\pm}$  was from  $C_{\pm} = 0$  until the value  $C_{\pm}^{(0)}$  where the conditions of Lemma 1 held. The value  $C_{\pm}^{(0)}$  was found for all  $\omega$  which we considered. This implies that all nonlinear modes within this range of  $\omega$  were found.

Fig.3 represents  $(\psi(0), \psi'(0))$ - picture for  $\omega = -3$ . Fig.4(a)-(b) shows four branches of localized modes  $\Gamma_r^n$ ,  $n = 1, 2, 3, 4$  which depart from the eigenvalues of linear problem (in this case no analytical form of linear modes is available). In addition, the branch  $\Gamma_r^2$  undergoes secondary *pitchfork bifurcation* and two branches  $\Gamma_r^{\pm}$  of nonsymmetric solutions arise. These solutions are localized in one of the wells of the potential  $V(x)$ . The linear stability analysis shows that the modes corresponding to the branches  $\Gamma_r^1$ ,  $\Gamma_r^3$ ,  $\Gamma_r^4$  as well as  $\Gamma_r^{\pm}$  are stable. The nonlinear modes corresponding to the branch  $\Gamma_r^2$  are stable until the point of pitchfork bifurcation. After the point of bifurcation they become unstable. This is quite expectable due to the results of the papers [25, 26, 27] and refines the statement of [12] where it has been suggested that all nonlinear modes in the case of repulsive nonlinearity are stable.

*3.2.2. Attractive nonlinearity.* In this case Eq.(8) reads

$$\psi'' + (\omega - \alpha x^4 - \beta x^2)\psi + \psi^3 = 0, \quad \alpha > 0, \quad \beta < 0 \quad (26)$$

The most part of calculations were done for  $\alpha = 10$ ,  $\beta = -20$ . Fig.4 represents all the branches of localized modes which were found within the range  $0 \leq C_{\pm} \leq 0.4$  and  $-20 \leq \omega \leq -4$ . There are two branches which correspond to modes with linear counterpart,  $\Gamma_a^1$  and  $\Gamma_a^2$  originated from 1-st and 2-nd eigenvalues of linear problem. At some value of  $\omega$  the branch  $\Gamma_a^1$  undergoes pitchfork bifurcation and two more branches of non-symmetric solution  $\Gamma_a^{\pm}$  emerge. Apart from them there is a branch  $\Gamma_a^0$  of nonlinear modes localized on the maximum of the potential situated between the two wells. These




**Figure 3.**  $(\psi(0), \psi'(0))$ -pictures for Eq.(8) with  $V(x) = 10x^4 - 20x^2$  and repulsive nonlinearity,  $\omega = -3$ .

solutions were reported also in [12]. When widening the window of scanning with respect to  $C_{\pm}$  and  $\omega$  much more branches appear; as a result, the complete picture becomes quite complex. The linear stability analysis shows that the solutions of the branch  $\Gamma_a^2$  as well as the branches  $\Gamma_a^{\pm}$  are stable, whereas the solutions of the branch  $\Gamma_a^0$  are unstable. As it was in repulsive case, the nonlinear modes corresponding to the branch  $\Gamma_a^1$  are stable until the point of pitchfork bifurcation and are unstable after this point. The part of solutions of the branch  $\Gamma_a^3$  shown in Fig.4(c)-(d) are unstable.

The presence of pitchfork bifurcation reflects the fact that Eq.(26) with even potential is invariant with respect to transformation  $x \rightarrow -x$ . The two nonsymmetric states  $\psi^{(+)}(x) \in \Gamma_a^+$  and  $\psi^{(-)}(x) \in \Gamma_a^-$  which bifurcate from the principal state  $\psi_1(x) \in \Gamma_a^1$  are connected by the symmetry  $\psi^{+}(-x) = \psi^{-}(x)$ . Adding small perturbation which breaks evenness of the potential results in replacing the pitchfork bifurcation by saddle-node one (see Fig.5). Similar phenomenon was reported in the paper [26] for the case of the potential composed by harmonic and periodic parts.

One more interesting phenomenon takes place when changing the parameters of the potential. It was found that if  $\alpha$  is fixed there exists a critical value of  $\beta$  when the branches  $\Gamma_a^1$  and  $\Gamma_a^0$  touch. When  $\beta$  passes through this critical value a reconnection of branches  $\Gamma_a^1$  and  $\Gamma_a^0$  occurs (see Fig.6). This means that by small variation of parameter  $\beta$  (keeping  $\alpha$  fixed) in vicinity of this critical point, a solution *with* linear counterpart




**Figure 4.** Panels (a)-(b): The branches of nonlinear modes for Eq.(8) with  $\alpha = 10$ ,  $\beta = -20$  and repulsive nonlinearity. (a):  $(C_+, \omega)$ -diagram, (b):  $(N, \omega)$ -diagram. The branches  $\Gamma_r^n$ ,  $n = 1, 2, 3, 4$  are originated by the modes of linear eigenvalue problem, the branches  $\Gamma_r^\pm$  appear as a result of secondary branching of  $\Gamma_r^2$  (the bifurcation is of pitchfork type). In  $(N, \omega)$ -diagram the branches  $\Gamma_r^+$  and  $\Gamma_r^-$  coincide.

Panels (c)-(d): The branches of nonlinear modes for Eq.(8) with  $\alpha = 10$ ,  $\beta = -20$  and attractive nonlinearity. (c):  $(C_+, \omega)$ -diagram, (d):  $(N, \omega)$ -diagram. The branches  $\Gamma_a^1$  and  $\Gamma_a^2$  are originated by the modes of linear eigenvalue problem, the branches  $\Gamma_a^\pm$  appear as a result of secondary branching of  $\Gamma_a^1$  (the bifurcation is of pitchfork type). In  $(N, \omega)$ -diagram the branches  $\Gamma_a^+$  and  $\Gamma_a^-$  coincide.

can be transformed into a solution *without* linear counterpart and vice versa.

#### 4. Conclusion

In the paper a method to study nonlinear modes for the Gross-Pitaevskii equation is described. It is based on exact statement on one-to-one correspondence between the solutions of Eq.(8) which vanish at  $x \rightarrow +\infty$  and a linear coefficient  $C_+$  which describe its asymptotic in this limit. This method can be regarded as an extension of shooting method used for calculation of single nonlinear mode (see e.g. [6, 20]) aimed to get a *complete picture* of coexisting nonlinear modes. Combined with some additional exact statements which have been proved in the case of repulsive nonlinearity the method




**Figure 5.**  $(C_+, \omega)$ -diagrams for Eq.(8), attractive case,  $V(x) = 10x^4 - 20x^2$  (left panel) and  $V(x) = 10x^4 - 20x^2 + 0.01x^3$  (right panel). Breaking of the symmetry of the potential results in replacing of pitchfork bifurcation by saddle-node one.




**Figure 6.** Transformation of  $(C_+, \omega)$ -diagram when the parameters of the potential vary. Left panel:  $\alpha = 10$ ,  $\beta = -20$ , central panel:  $\alpha = 10$ ,  $\beta = -15.91$ , right panel:  $\alpha = 10$ ,  $\beta = -15.5$ . Only branches of even solutions are depicted.

allows to fulfill demonstrative computation to guarantee that *all* the nonlinear modes corresponding to given value of  $\omega$  (chemical potential in terms of BEC) are found.

Conforming to the approach which we describe we offer a representation of the branches of solutions on the plane  $(C_+, \omega)$  instead of more traditional representation on the plane  $(N, \omega)$ ,  $N = \int_{-\infty}^{\infty} \psi^2(x) dx$ . The advantage of this representation is that each point of the plane  $(C_+, \omega)$  corresponds to at most one nonlinear mode whereas the point on the plane  $(N, \omega)$  can correspond to many nonlinear modes which share the same number of particles. Therefore the intersection (or touching) of the curves which correspond to the branches of nonlinear modes in the plane  $(C_+, \omega)$  means that some bifurcation occurs. This is not the case for  $(N, \omega)$  representation. The  $(C_+, \omega)$ -diagrams calculated in the case of harmonic potential  $V(x) = x^2$  indicate that all the nonlinear modes have linear counterpart. In the case of double-well potential the situation is richer, due to modes without linear counterpart and secondary branching of non-symmetric modes.

In the paper the approach is applied to 1D GPE with potential having one or two minima. However it seems to be an interesting problem to apply this method for the study of 1D GPE with potentials with three or more minima (see e.g. [28]). Also it may

be used for GPE with periodic potential (see e.g. [29, 30]) as well as for the potential generated by magnetic fields and optical lattice [26]. Applications to quintic or cubic-quintic GPE model are also possible [31, 32]. With minor modifications this approach can be applied to 2D and 3D radial versions of GPE.

## Acknowledgments

Authors are grateful to Prof. V.Konotop and Prof. L.Lerman for fruitful discussions and pointing to useful references. G.A. acknowledges the support from President Program in Support for Leading Scientific Schools in Russia (Project 4122.2006.2.).

## Appendix A. Summary of results for the equation $\psi'' - Q(x)\psi = 0$ .

The equation

$$\psi'' - Q(x)\psi = 0 \quad (\text{A.1})$$

(the prime denotes the derivative with respect to  $x$ ) has been discussed in numerous books and papers [15, 16, 18]. Let us set forth below some results about Eq. (A.1) which have been used in this paper.

Statement A.1:(see [15], chapter 2) *Let there exists some  $x_0$  and  $\varepsilon$  such that*

$$Q(x) \geq \varepsilon > 0 \quad \text{for } x > x_0. \quad (\text{A.2})$$

*Then a pair of linearly independent solutions of (A.1),  $\psi_1(x)$  and  $\psi_2(x)$ , can be chosen in such a way that for some value  $a > x_0$  the following assertions hold:*

(A1) *the functions  $\psi_1(x)$ ,  $\psi_1'(x) \rightarrow 0$  when  $x \rightarrow +\infty$ ;*

(A2) *the functions  $\psi_1(x)$ ,  $\psi_1'(x)$  have no zeros and are monotonic for  $x > a$ ;*

(A3) *for  $x > a$*

$$|\psi_1(x)| \leq A_1 e^{-\sqrt{\varepsilon}x} \quad (\text{A.3})$$

*for some  $A_1$ .*

(A4) *the solution  $\psi_1(x)$  is unique (up to constant factor).*

(B1) *the functions  $\psi_2(x)$ ,  $\psi_2'(x) \rightarrow \infty$  when  $x \rightarrow +\infty$ ;*

(B2) *the functions  $\psi_2(x)$ ,  $\psi_2'(x)$  have no zeros and are monotonic for  $x > a$ ;*

(B3) *for  $x > a$*

$$|\psi_2(x)| \geq A_2 e^{\sqrt{\varepsilon}x} \quad (\text{A.4})$$

*for some  $A_2$ .*

Comment: The points (A1)-(A3) and the fact that the sign of  $\psi''(x) > 0$  coincides with the sign of  $\psi(x)$  for  $[a; +\infty)$  imply that

$$\psi_1(x), \psi_1'(x), \psi_1''(x) \in L_p[a; +\infty); \quad p > 0 \quad (\text{A.5})$$

Imposing more restrictions to  $Q(x)$  results in more precise description for the solutions  $\psi_1(x)$  and  $\psi_2(x)$  growth or decay:

Statement A.2:(see [22], Chapter IV, Theorem 14) *Let the function  $Q(x)$  satisfies the condition (A.2). Let, in addition, the following conditions hold:*

$$(a) \ Q(x) \in C^2(x_0, +\infty);$$

$$(b) \ \int_{x_0}^{\infty} \frac{|Q''(x)|}{Q^{3/2}(x)} dx < \infty,$$

*Then the solutions  $\psi_1(x)$ ,  $\psi_2(x)$  (which exist due to Statement A.1) obey the following asymptotic formulas:*

$$\psi_1(x) = \frac{e^{-S(x_0, x)}}{Q^{1/4}(x)} (1 + o(1)), \quad (A.6)$$

$$\psi_2(x) = \frac{e^{S(x_0, x)}}{Q^{1/4}(x)} (1 + o(1)), \quad S(x_0, x) = \int_{x_0}^x \sqrt{Q(t)} dt \quad (A.7)$$

Moreover, the asymptotic formulas (A.7)-(A.6) can be differentiated at least once.

Comment: Due to (A.2)  $S(x_0, x) \rightarrow +\infty$  as  $x \rightarrow +\infty$ .

Comment: Evidently, there is a freedom in choosing the lower bound of integration in (A.7)-(A.6). Its change from  $x = x_0$  to  $x = x_1$  results in rescaling of  $\psi_1(x)$  and  $\psi_2(x)$

$$\psi_1(x) \longrightarrow e^{S(x_0, x_1)} \psi_1(x); \quad \psi_2(x) \longrightarrow e^{-S(x_0, x_1)} \psi_2(x)$$

## Appendix B. Proof of Theorem from Section 2.2.

The aim of this Appendix is to prove the following statement for the equation

$$\psi'' - Q(x)\psi + F(\psi) = 0 \quad (B.1)$$

Theorem 1. *Let there exists some  $x_0$  such that*

$$(Q1) \ Q(x) \geq \varepsilon > 0 \text{ for } x > x_0;$$

$$(Q2) \ Q(x) \in C^2(x_0, +\infty);$$

$$(Q3) \ \int_{x_0}^{\infty} \frac{|Q''(x)|}{Q^{3/2}(x)} dx < \infty$$

*and there exist some  $\delta$  such that*

$$(NL1) \ F(t) \in C^3[-\delta; \delta].$$

$$(NL2) \ F(0) = F'(0) = 0;$$

*Let  $\psi_1(x)$  be the solution of linear equation  $\psi'' - Q(x)\psi = 0$  such that  $\psi_1(x) \rightarrow 0$  as  $x \rightarrow +\infty$  (see Appendix A). Let  $S_+$  be a set of all solutions of Eq.(B.1) such that  $\psi(x) \rightarrow 0$  when  $x \rightarrow +\infty$ . Then*

*(a) for any solution  $\psi(x) \in S_+$  there exists a limit*

$$\lim_{x \rightarrow +\infty} \frac{\psi(x)}{\psi_1(x)} = C < \infty \quad (B.2)$$

(b) for any  $C \in \mathbf{R}$  there exists an unique solution  $\psi_+(x; C) \in S_+$  of Eq.(B.1) such that

$$\lim_{x \rightarrow +\infty} \frac{\psi_+(x, C)}{\psi_1(x)} = C \quad (\text{B.3})$$

(c) if  $\psi_+(x; C_+)$  exists in some neighbourhood  $U$  of a point  $(\tilde{x}; \tilde{C}_+)$  then in  $U$  there also exists partial derivative  $\frac{\partial \psi_+(x; C_+)}{\partial C_+}$  which is continuous with respect to  $C_+$ .

Comment: If for a function  $F(t)$  the condition (NL2) holds and  $F(t) \in C^2[-\delta; \delta]$  for some  $\delta > 0$  then  $f(t) \equiv F(t)/t$  has no singularities in  $|t| < \delta$  and there exists  $A_0 > 0$  and  $\delta_1$  such that

$$|f(t)| \leq A_0|t|, \quad |t| < \delta_1; \quad (\text{B.4})$$

The proof of Theorem 1 can be sketched in a form of a sequence of Lemmas. First Lemma states that small perturbations of potential in Eq.(A.1) do not change the asymptotic (A.6) of the solution which tends to zero as  $x \rightarrow +\infty$ .

Lemma B.1 Let  $Q(x)$  satisfy the conditions (Q1)-(Q3) of Theorem 1 for  $x > x_0$  and  $\psi_1(x)$  be a solution of equation  $\psi'' - Q(x)\psi = 0$  such that  $\psi_1(x) \rightarrow 0$  as  $x \rightarrow +\infty$  (guaranteed by Statement A.1). Let  $R(x) \in C^2(x_0, +\infty)$ ,  $R(x) \rightarrow 0$  when  $x \rightarrow +\infty$  and  $R(x), R''(x) \in L_1(x_0, +\infty)$ . Then the equation

$$\eta'' - (Q(x) + R(x))\eta = 0 \quad (\text{B.5})$$

has unique (up to constant factor) solution  $\eta_1(x)$  such that:

- (i)  $\eta_1(x), \eta_1'(x)$  are monotonous functions on some interval  $[a; +\infty)$ ,  $a \geq x_0$ ;
- (ii)  $\eta_1(x), \eta_1'(x)$  tend to zero when  $x \rightarrow +\infty$ ;
- (iii) the solution  $\eta_1(x)$  when  $x \rightarrow +\infty$  has the asymptotic

$$\eta_1(x) = \psi_1(x)(C_1 + o(1)); \quad \eta_1'(x) = \psi_1'(x)(C_1 + o(1)) \quad (\text{B.6})$$

for some  $C_1 \in \mathbf{R}$ .

Proof: It is straightforward to check that  $Q_1(x) \equiv Q(x) + R(x)$  also satisfies the conditions of Statement A.2. Then the statement of Lemma B.1 follows immediately from formula (A.6). ■

Lemma B.2 Let  $Q(x)$  satisfies the conditions (Q1)-(Q3) of Theorem 1. Let  $F(t) \in C^2[-\delta, \delta]$  for some  $\delta > 0$  and satisfies the condition (NL2). Then for any solution  $\psi(x)$  of Eq.(B.1) such that  $\psi(x) \rightarrow 0$  as  $x \rightarrow +\infty$  there exists some  $C \in \mathbf{R}$  such that the asymptotic of  $\psi(x)$  is given by formula

$$\psi(x) = \frac{e^{-S(x_0, x)}}{Q^{1/4}(x)} (C + o(1)), \quad S(x_0, x) = \int_{x_0}^x \sqrt{Q(t)} dt \quad x \rightarrow +\infty \quad (\text{B.7})$$

The case  $C = 0$  corresponds to zero solution. This asymptotic formula can be differentiated at least once.

Proof: Due to the condition (NL2) one can introduce  $f(t) = F(t)/t$  such that  $f(0) = 0$ . Suppose that  $\psi(x)$  is a solution of Eq.(B.1) such that  $\psi(x) \rightarrow 0$  as  $x \rightarrow +\infty$ . Let us rewrite Eq.(B.1) in the form

$$\psi'' - (Q(x) + R(x))\psi = 0, \quad R(x) = -f(\psi(x));$$

One can show that the conditions of Lemma B.2 imply the conditions of Lemma B.1 for  $R(x)$ . Then the statement of Lemma B.2 follows from Lemma B.1. ■

Lemma B.3 *Let  $Q(x)$  satisfies the conditions (Q1)-(Q3) for some  $x > x_0$ . Let  $F(t) \in C[-\delta, \delta]$  for some  $\delta > 0$  and  $F(t)$  satisfies the condition (B.4) for  $|t| < \delta$ . Let  $\psi_1(x)$  be some fixed nonzero solution of linear problem (A.1), such that  $\psi_1(x) \rightarrow 0$  when  $x \rightarrow +\infty$ . Then for any  $C \in \mathbf{R}$  there exists a solution  $\psi_+(x; C)$  of (B.1) such that*

$$\lim_{x \rightarrow +\infty} \frac{\psi_+(x; C)}{\psi_1(x)} = C \quad (\text{B.8})$$

Comment. This statement is converse to Lemma B.2. The assertion of Lemma B.3 says that for any  $C$  there exists a solution of (B.1) which at  $x \rightarrow +\infty$  has the asymptotic

$$\psi_+(x) = \frac{e^{-S(x_0, x)}}{Q^{1/4}(x)} (C + o(1)), \quad S(x_0, x) = \int_{x_0}^x \sqrt{Q(t)} dt \quad (\text{B.9})$$

Proof: Let  $\psi(x)$  satisfy the equation (B.1). Introduce the function

$$\Psi(x) = \frac{\psi(x)}{\psi_1(x)} - C$$

where  $C$  is free real parameter and  $\psi_1(x)$  is fixed solution of linear equation (A.1), such that  $\psi_1(x) \rightarrow 0$  as  $x \rightarrow +\infty$ . Then  $\Psi(x)$  satisfies the system of equations

$$\Psi' = \Phi \quad (\text{B.10})$$

$$\Phi' = -P(x)\Phi - (\Psi + C) \cdot f((\Psi + C)\psi_1(x)), \quad P(x) = \frac{2\psi_1'(x)}{\psi_1(x)} \quad (\text{B.11})$$

where  $f(t) = F(t)/t$ . The existence of solution  $\Psi(x)$  such that  $\Psi(x) \rightarrow 0$  as  $x \rightarrow +\infty$  follows from Theorem 9.1, Chapter XII of [19] applied to the system (B.10)-(B.11). The conditions of this Theorem can be verified straightforwardly ■.

The following Lemma says that for given  $C$  the solution of (B.1) with constant  $C$  in asymptotic (B.7) is unique.

Lemma B.4 *Let  $Q(x)$  satisfies the conditions (Q1)-(Q3) of Theorem 1 for  $x > x_0$  and  $F(t)$  satisfies the conditions (NL1)-(NL2). Then for any two solutions,  $\psi_a(x)$  and  $\psi_b(x)$ , of Eq.(B.1) such that  $\psi_a(x), \psi_b(x) \rightarrow 0$  when  $x \rightarrow \infty$  and  $\psi_a(x) \not\equiv \psi_b(x)$  the following inequality holds:*

$$\lim_{x \rightarrow +\infty} \frac{\psi_a(x)}{\psi_b(x)} \neq 1 \quad (\text{B.12})$$

*Proof:* Assume the contrary, that  $\lim_{x \rightarrow +\infty} \frac{\psi_a(x)}{\psi_b(x)} = 1$ . Then, according to Lemma B.2, both solutions  $\psi_a(x)$  and  $\psi_b(x)$  share the same asymptotics

$$\psi_a(x) = \frac{e^{-S(x_0, x)}}{Q^{1/4}(x)} (C + o(1)), \quad \psi_b(x) = \frac{e^{-S(x_0, x)}}{Q^{1/4}(x)} (C + o(1)), \quad (\text{B.13})$$

$$S(x_0, x) = \int_{x_0}^x \sqrt{Q(t)} dt \quad (\text{B.14})$$

Subtracting the second equation from the first one in the system

$$\psi_b'' - Q(x)\psi_b + F(\psi_b) = 0, \quad \psi_a'' - Q(x)\psi_a + F(\psi_a) = 0$$

we conclude that the difference  $\Delta = \psi_b - \psi_a$  obeys the equation

$$\Delta'' - (Q(x) + G(\psi_a(x), \psi_b(x))) \Delta = 0, \quad (\text{B.15})$$

$$G(\psi_a(x), \psi_b(x)) = -\frac{F(\psi_b(x)) - F(\psi_a(x))}{\psi_b(x) - \psi_a(x)} \quad (\text{B.16})$$

Let us denote  $R(x) = G(\psi_a(x), \psi_b(x))$ . It is tedious but straightforward to show that  $R(x)$  satisfies the conditions of Lemma B.1 (the proof involves Lagrange theorem). Then it follows from Lemma B.1 that

$$\Delta(x) = \frac{e^{-S(x_1, x)}}{Q^{1/4}(x)} (C_1 + o(1)), \quad S(x_1, x) = \int_{x_1}^x \sqrt{Q(t)} dt \quad (\text{B.17})$$

Since  $\Delta \neq 0$  then  $C_1 \neq 0$  and  $\lim_{x \rightarrow +\infty} \frac{\Delta(x)}{\psi_b(x)} = \frac{C_1}{C} \neq 0$  that proves Lemma B.4. ■

Combining the results of Lemmas B.2, B.3, B.4 we arrive at the statements (a) and (b) of Theorem 1. The statement (c) follows from the uniqueness of the solution  $\psi_+(x; C)$  for given  $C$  (the statement (b)) and the property (NL1) (see e.g. [19]). This completes the proof of Theorem 1. ■

## Appendix C. Proof of Lemmas 1 and 2 from Section 2.4

Comparison Lemma. Let the functions  $y(t)$  and  $x(t)$ ,  $t \in [a; b]$  be solutions of equations

$$y_{tt} - g(t, y) = 0 \quad (\text{C.1})$$

$$x_{tt} - f(t, x) = 0 \quad (\text{C.2})$$

correspondingly. Let also the following conditions hold:

(i)  $f(t, \xi)$ ,  $g(t, \xi)$  are defined on  $[a; b] \times [A; B]$  and locally Lipschitz continuous (see [24]) with respect to  $\xi$ ,  $\xi \in [A; B]$ , ( $A, B$ ,  $b$  may be finite or infinite);

(ii)  $g(t, \xi) \geq f(t, \xi)$  for any  $t \in [a; b]$ ,  $\xi \in [A; B]$ ;

(iii)  $f(t, \xi)$  is monotone nondecreasing with respect to  $\xi$ ,  $\xi \in [A; B]$ .

Let  $A < x(a) \leq y(a) < B$  and  $x_t(a) \leq y_t(a)$ . Then  $x_t(t) \leq y_t(t)$  and  $x(t) \leq y(t)$  while  $A < x(t), y(t) < B$  or for the whole interval  $t \in [a; b]$

*Proof:* Introduce the functions

$$\tilde{f}(t, \xi) = \begin{cases} f(t, A), & \xi < A \\ f(t, \xi), & A \leq \xi \leq B \\ f(t, B), & \xi > B \end{cases} \quad \tilde{g}(t, \xi) = \begin{cases} g(t, A), & \xi < A \\ g(t, \xi), & A \leq \xi \leq B \\ g(t, B), & \xi > B \end{cases}$$

and let  $\tilde{x}(t), \tilde{y}(t)$  be the solutions of Cauchy problem for the equations

$$\tilde{y}_{tt} - \tilde{g}(t, \tilde{y}) = 0$$

$$\tilde{x}_{tt} - \tilde{f}(t, \tilde{x}) = 0$$

such that  $\tilde{x}(a) = x(a)$ ,  $\tilde{x}_t(a) = x_t(a)$ ,  $\tilde{y}(a) = y(a)$ ,  $\tilde{y}_t(a) = y_t(a)$ . The functions  $\tilde{x}(t)$  and  $\tilde{y}(t)$  coincide with  $x(t)$ ,  $y(t)$  when  $A < x(t), y(t) < B$ . Consider vectors  $\mathbf{v}, \mathbf{u}$  and vector-functions  $\mathbf{G}(t; \mathbf{v})$ ,  $\mathbf{F}(t, \mathbf{u})$  defined as

$$\mathbf{v} = \begin{pmatrix} \tilde{y} \\ \tilde{y}_t \end{pmatrix}; \quad \mathbf{G}(t, \mathbf{v}) = \begin{pmatrix} \tilde{y}_t \\ \tilde{g}(t, \tilde{y}) \end{pmatrix};$$

$$\mathbf{u} = \begin{pmatrix} \tilde{x} \\ \tilde{x}_t \end{pmatrix}; \quad \mathbf{F}(t, \mathbf{u}) = \begin{pmatrix} \tilde{x}_t \\ \tilde{f}(t, \tilde{x}) \end{pmatrix}$$

For  $\mathbf{u}_1, \mathbf{u}_2 \in \mathbf{R}^2$  we define the order relation:  $\mathbf{u}_1 \leq \mathbf{u}_2$  if each component of  $\mathbf{u}_2$  is greater or equal than the corresponding component of  $\mathbf{u}_1$ . The function  $\mathbf{F}(t, \mathbf{u})$  is monotone nondecreasing in the sense that if  $\mathbf{u}_1 \leq \mathbf{u}_2$  then  $\mathbf{F}(t, \mathbf{u}_1) \leq \mathbf{F}(t, \mathbf{u}_2)$ . Then

$$\mathbf{0} = \mathbf{u}_t - \mathbf{F}(t, \mathbf{u}) = \mathbf{v}_t - \mathbf{G}(t, \mathbf{v}) \leq \mathbf{v}_t - \mathbf{F}(t, \mathbf{v})$$

In this situation Theorem 2 of [23] (see also [24]) states that  $\mathbf{u}(a) \leq \mathbf{v}(a)$  implies that  $\mathbf{u}(t) \leq \mathbf{v}(t)$  for  $t \in [a; b]$ . This gives the assertion of Comparison Lemma. ■

*Proof of Lemma 1.* Let us denote  $u(x, C) = \frac{\partial \psi_+(x; C)}{\partial C}$ . Then  $u(x, C)$  satisfies the linear equation

$$u'' + (\omega - V(x) - 3\psi_+^2(x, C))u = 0 \quad (\text{C.3})$$

The conditions of Lemma 1 imply that there is an interval  $[C_+^0, C_+^1]$  such that for any  $C \in [C_+^0, C_+^1]$  and any  $x$  for which the solution  $\psi_+(x, C)$  is defined, the following conditions hold

$$\psi_+(x; C) > 0 \quad (\text{C.4})$$

$$\omega - V(x) - 3\psi_+^2(x; C) < 0 \quad (\text{C.5})$$

Let at least one of the conditions (C.4)-(C.5) breaks at  $C = C_+^1$  i.e. one at least one of the signs “<”, “>” in (C.4)-(C.5) becomes “=” at some value  $x = a$ . It follows from asymptotics (15) that there exists  $b$  large enough that  $\psi_+(x; C)$  for  $x > b$  and  $C \in [C_+^0, C_+^1]$  is monotonic with respect to  $C$  in the sense that if  $C_1 \geq C_2$ ,  $C_{1,2} \in [C_+^0, C_+^1]$  and  $x > b$  then  $\psi_+(x; C_1) \geq \psi_+(x; C_2)$  and  $\psi'_+(x; C_1) \leq \psi'_+(x; C_2)$ . This means that for  $x > b$  and  $C \in [C_+^0, C_+^1]$  one has  $u(x; C) \geq 0$  and  $u'(x; C) \leq 0$ . After changing  $x \rightarrow -x$  one can apply Comparison Lemma to equations (C.3) and  $u'' = 0$ . Then  $u(x; C) \geq 0$  for  $C \in [C_+^0, C_+^1]$  not only when  $x > b$  but everywhere in the interval where  $u(x; C)$  is

defined. This implies that  $\frac{\partial \psi_+(a;C)}{\partial C} \geq 0$  for  $C \in [C_+^0, C_+^1]$  i.e.  $\psi_+(a;C)$  is nondecreasing function of  $C$ . Therefore  $\psi_+(a;C_+^0) \leq \psi_+(a;C_+^1)$  and this is a contradiction to the assumption that one of the conditions (C.4)-(C.5) breaks at  $C = C_+^1$ . ■

*Proof of Lemma 2.* Let us apply Comparison Lemma taking

$$f(t, \xi) = -\Omega\xi + \xi^3; \quad g(t, \xi) = -(\omega - V(t))\xi + \xi^3 \quad (\text{C.6})$$

$$A = \sqrt{\Omega}; \quad B = +\infty \quad (\text{C.7})$$

It is straightforward to check that any solution of the equation  $\psi'' + \Omega\psi - \psi^3 = 0$  with initial data  $\psi(t_0) \geq \sqrt{\Omega}$ ,  $\psi'(t_0)$  - arbitrary, is singular. Therefore any solution of Eq(20) with the same initial conditions is also singular. Due to oddness of the function  $g(t, \xi)$  with respect to  $\xi$  one concludes that any solution of Eq(20) such that  $\psi(t_0) \leq -\sqrt{\Omega}$ ,  $\psi'(t_0)$  - arbitrary, also is singular. This implies the assertion of Lemma 2. ■

## References

- [1] F.Dalfovo, S.Giorgini, L.P.Pitaevskii, Rev.Mod.Phys., **71**, 463, (1999)
- [2] M.H.Anderson, J.R.Ensher, M.R.Matthews, C.E.Wieman and E.A.Cornell, Science, **269**, 198 (1995)
- [3] C.C.Bradley, C.A.Sackett, J.J.Tollett and R.G.Hulet, Rhys. Rev. Letters, **75**, 1687, (1995)
- [4] M.Edwards, K.Burnett, Phys.Rev.A, **51**,(2), 1382 (1995)
- [5] F.Dalfovo, S.Stringari, Phys.Rev.A, **53**(4), 2477 (1996)
- [6] P.A.Ruprecht, M.J.Holland, K.Burnett, M.Edwards, Phys.Rev.A, **51**, (6), 4704,(1995)
- [7] V.I. Yukalov, E.P. Yukalova and V.S. Bagnato, Phys. Rev. A **56**, 4845 (1997), Phys. Rev. A **66**, 043602 (2002)
- [8] V.I. Yukalov, E.P. Yukalova and V.S. Bagnato, Phys. Rev. A **66**, 043602 (2002)
- [9] Yu.S.Kivshar, T.J.Alexander and S.K.Turitsyn, Phys.Lett.A, **278**,225, (2001)
- [10] M. Kunze, T. Kupper, V. K. Mezentsev, E. G. Shapiro, and S. K. Turitsyn, Phys. D, **128**, 273, (1999).
- [11] R.D'Agosta, B.A.Malomed and C.Presilla. Laser Physics, **12**, (1), 37 (2002)
- [12] R. D'Agosta and C. Presilla, Phys. Rev. A, **65**, 043609, (2002)
- [13] M. Brtko, A. Gammal, L. Tomio, Phys. Lett A **359**, 339, (2006)
- [14] P. Zi, E. Infeld, M. Matuszewski, G. Rowlands and M. Trippenbach, Phys. Rev. A., **73**, 022105, (2006)
- [15] F.A.Berezin, M.A. Shubin. The Shrödinder equation. Kluwer Academic Publisher,1991
- [16] M.V. Fedoryuk, Asymptotic Analysis. Linear Ordinary Differential Equations, Springer-Verlag, Berlin, 1993; see also M.V. Fedoryuk, Asymptotics of solutions of linear differential equations of the second order in complex domain, Addendum 1 to russian translation of: Wasov, W., Asymptotic Expansions for Ordinary Differential Equations, Moscow, Mir, 1968 (in Russian).
- [17] H.Leiva, H.M.Rodrigues, Nonlinear Analysis, **47**, 4579 (2001)
- [18] E. Kamke, Differentialgleichungen: Lösungsmethoden und Lösungen, I, Gewöhnliche Differentialgleichungen, B. G. Teubner, Leipzig, 1977
- [19] Ph.Hartman, Ordinary Differential Equations, John Wiley and Sons, New York, 1964.
- [20] A.Gammal, T.Frederico and L. Tomio, Phys.Rev. E, **60**, 2421, (1999)
- [21] P.G.Kevrekidis, V.V.Konotop, A.Rodrigues and D.J. Frantzeskakis, J.Phys.B:At.Mol.Opt.Phys. **38**, 1173 (2005)
- [22] W.A.Coppel. Stability and asymptotic behaviour of differential equations, Heath and Co, Boston, 1965.
- [23] G.Herzog. Nonlinear Analysis, **47**, 2213, (2001)

- [24] G.Herzog, R.Lemmert. Differential inequalities, <http://www.mathematik.uni-karlsruhe.de/~semlv>
- [25] R.K. Jackson and M.I.Weinstein. J. Stat. Phys., **116**, 881, (2004).
- [26] T. Kapitula and P.G. Kevrekidis. Nonlinearity, **18**, 2491, (2005).
- [27] P.G. Kevrekidis, Zh. Chen, B.A. Malomed, D.J. Frantzeskakis and M.I. Weinstein. Phys. Lett. A, **340**, 275, (2005).
- [28] T. Kapitula, P. Kevrekidis, and Z. Chen, SIAM J. Appl. Dyn. Sys. 5(4), p.598 (2006)
- [29] G.L. Alfimov, V.V. Konotop, and M. Salerno, Europhys. Lett. 58, 7, (2002).
- [30] P. J. Y. Louis, E. A. Ostrovskaya, C. M. Savage, and Yu. S. Kivshar, Phys. Rev. A 67, 013602, (2003)
- [31] F.Abdullaev, M.Salerno, Phys. Rev. A 72, 033617 (2005).
- [32] G.Alfimov, V.Konotop and P.Pacciani, Phys.Rev.A, to appear.